COMMENT

Sources of Dietary Iodine: Bread, Cows' Milk, and Infant Formula in the Boston Area

ELIZABETH N. PEARCE, SAM PINO, XUEMEI HE, HAMID R. BAZRAFSHAN, STEPHANIE L. LEE, AND LEWIS E. BRAVERMAN

Section of Endocrinology, Diabetes, and Nutrition (E.N.P., S.P., X.H., S.L.L., L.E.B.), Boston Medical Center, Boston University School of Medicine, Boston, Massachusetts 02118; and Golestan University Medical School (H.R.B.), Golestan, Iran

Dietary iodine is essential for thyroid hormone production. Although U.S. dietary iodine is generally adequate, some groups, especially women of childbearing age, are at risk for mild iodine deficiency. Children's average urinary iodine is higher than that of adults. U.S. dietary iodine sources have not been assessed recently. A survey of iodine content in 20 brands of bread, 18 brands of cows' milk, and eight infant formulae was performed between 2001 and 2002. Three bread varieties contained more than 300 μ g iodine per slice. Iodine content in

other brands was far lower (mean \pm SD, $10.1 \pm 13.2~\mu g$ iodine/ slice). All cows' milk samples had at least 88 μg iodine/250 ml, ranging from 88–168 μg (116.0 \pm 22.1 μg /250 ml). Infant formulae values ranged from 16.2 to 56.8 μg iodine/5 oz (23.5 \pm 13.78 μg /5 oz). The public should be aware of the need for adequate dietary iodine intake and should be aware that ingredient lists do not reflect the iodine content of foods. (*J Clin Endocrinol Metab* 89: 3421–3424, 2004)

DEQUATE DIETARY IODINE intake is essential for the production of thyroid hormones. The recommended daily allowances for iodine vary by age (Table 1) (1). Iodine intake in the United States has varied over the years due to alterations in the iodine content of foodstuffs. Since the iodization of salt and other foods in the 1920s, U.S. dietary iodine has generally been adequate. Subjects sampled for the most recent U.S. National Health and Nutrition Examination Survey (NHANES 2000) had a median urinary iodine excretion of 16.1 μ g/dl (National Center for Health Statistics: Health E-Stats: Iodine Level, United States, 2000; http:// www.cdc.gov/nchs/products/pubs/pubd/hestats/iodine. htm), which was essentially unchanged from the previous survey in 1988–1994 (NHANES III) (2). However, NHANES III demonstrated an approximate 50% reduction in median dietary iodine intake (median urinary iodine, 14.5 µg/dl) compared with NHANES I values from 1971-1974 (median urinary iodine, 32.0 μ g/dl) (2). In particular, some subsets of the population, especially women of childbearing age, had an increased prevalence of mild iodine deficiency. This is of concern because thyroid hormone, requiring adequate iodine intake, is critical for neural development *in utero* and in early life. Although cretinism due to iodine deficiency is not a problem in the United States, subtle developmental delays could result from mild maternal iodine deficiency. Another population subset, school-aged children, had urinary iodine

JCEM is published monthly by The Endocrine Society (http://www.endo-society.org), the foremost professional society serving the endocrine community.

excretion in both the NHANES I and III surveys that was higher than that of adults (2).

In the United States, iodine deficiency has been largely eliminated by means of silent prophylaxis. Iodine supplementation of salt and other foods has never been mandated, and iodine content of most foods is not listed on package labels. It has been over 10 yr since the last market basket analysis for U.S. iodine intake was performed (3). Therefore, it has been difficult to determine the primary sources of iodine in the U.S. diet. In particular, the higher median urinary iodine levels in children compared with adults, reflecting dietary iodine intake, have not been explained.

Materials and Methods

To document some potentially important sources of iodine in the U.S. diet, we measured iodine content in a variety of locally available dietary staples during 2001–2002.

Two slices of bread were obtained from 20 different brands of sliced bread from supermarkets in the Boston area. An initial sample of breads was brought from home by co-workers; several more brands were then selected randomly at a local supermarket. Iodine content was analyzed in 100–150 mg of each slice and measured in duplicate. Seven brands, all products of the Interstate Bakeries Corp. (Charlotte, NC), listed iodate conditioners as ingredients; the iodine content was not indicated. None of the store-bought breads listed iodized salt as an ingredient. When some of the brands that listed iodate conditioners in the ingredients list were tested, the samples contained only small amounts of iodine. The manufacturer was contacted and indicated that the iodine conditioners had been removed from all the breads at the last recipe reformulation and that supplies of the old bread bags were being depleted.

The iodine content of 18 brands of milk sold in local supermarkets was measured. Because cows' milk from some processing plants is sold under multiple brand names, we effectively measured milk from five separate sources. Measurements were performed twice on each brand, once in February and once in August, to determine whether there was any seasonal variation in milk iodine content. Milk iodine content was

Abbreviation: NHANES, National Health and Nutrition Examination

not included on any of the labels. The iodine content of eight infant formulae, representing all types available at a single large local supermarket, was also measured and compared with their labeled iodine

Total iodine concentrations in cows' milk, infant formulae, and bread were measured spectrophotometrically by a modification of the method of Benotti et al. (4). All samples were digested with 3.0 ml chloric acid (30% aqueous solution) before analysis. Iodine concentrations from all samples were measured at least twice; in 95% of the samples, the initial two measurements were within 15% of each other, and the two values were averaged. In cases where the initial two measurements were not within 15% of each other, a third or a fourth measurement was obtained, and the average of all measurements was reported.

Statistical analyses were performed using SAS version 8 (SAS Institute, Cary, NC). Paired t tests were used to compare mean iodine content in winter compared with summer cows' milk samples.

TABLE 1. United States Institute of Medicine recommendations for iodine intake

Group	Iodine intake (μg/d)		
Age 0-6 months	$110 (AI)^a$		
Age 7–12 months	130 (AI)		
Age 1–8 yr	$90~(\mathrm{RDA})^a$		
Age 9–13 yr	120 (RDA)		
Age 14 yr and over	150 (RDA)		
Pregnant women	220 (RDA)		
Lactating women	290 (RDA)		

AI, Adequate intake; RDA, recommended daily allowance.

 a The RDA is the intake of a nutrient expected to meet the needs of 97–98% of healthy individuals. The AI is an approximation of the dietary intake of healthy people who are assumed to be getting adequate nutrition. The AI is used when there is not enough evidence to determine the RDA; it always exceeds the RDA.

Results

Three varieties of bread contained more than 313 µg iodine per slice (range, 313.5–587.4 μ g). Iodine content in the other 17 brands ranged from 2.2-54 µg iodine per slice (mean, $10.1 \pm 13.2 \,\mu g$ iodine/slice; Table 2).

All 18 brands of cows' milk had at least 88 µg iodine per 250 ml (\sim 8 oz or 1 cup), ranging from 88–168 μ g (mean \pm sp, $116.0 \pm 22.1 \,\mu\text{g}/250$ ml; Fig. 1). Mean iodine content of




Fig. 1. Iodine concentrations in summer and winter cows' milk samples by source dairy.

TABLE 2. Iodine concentration in breads from the Boston area

Bread brand	Weight per slice (g)	$\begin{array}{c} \text{Iodine} \\ (\mu \text{g/g}) \end{array}$	Total iodine per slice (μg)
Arnold Jewish Bread ^a	31.5	0.11	3.5
Bouyea Fassett 12 Grain Bread ^a	47.4	0.10	4.6
Cape Cod Wheat Bread	33.6	0.35	11.8
Country Kitchen Butter Split Wheat ^b	28.4	0.09	2.6
Country Kitchen Giant White ^b	30.0	0.13	3.8
Country Kitchen Hearty Canadian Brown ^b	34.4	0.09	3.2
Finagle a Bagel Rye Bagel ^c	132.1	0.11	13.9
Home Pride Butter Top Wheat d	29.0	0.19	5.4
Home Pride Butter Top White ^d	30.6	0.21	6.5
J.J. Nissen Italian ^d	28.2	0.11	3.1
J.J. Nissen Split Top Wheat ^d	30.0	0.19	5.7
Martin's Potato Bread ^e	33.9	0.49	16.6
Pepperidge Whole Grain ^f	29.0	0.12	3.5
Shaw's Foccacia ^g	25.2	23.31	587.4
Shaw's Giant White ^g	25.8	1.13	29.2
Shaw's Marble Rye ^g	25.3	2.12	53.5
Sunbeam Batter Whipped White ^d	26.3	12.49	328.0
Sunbeam White ^d	26.7	12.15	324.3
When Pigs Fly Six Grain and Pumpkin Seed ^h	51.2	0.06	2.8
Wonder Bread^d	25.4	0.09	2.2

^a Products of George Weston Bakeries, Inc. (Totowa, NJ).

^b Products of LePage Bakeries (Auburn, ME).

^c Product of Finagle a Bagel (Boston, MA).

^d Products of the Interstate Bakeries Corp. (Charlotte, NC).

^e Product of Martin's Famous Pastry Shoppe, Inc. (Chambersburg, PA).

f Product of Campbell Soup Co. (Camden, NJ).
g Products of Shaw's Supermarkets, Inc. (West Bridgewater, MA).

^h Product of When Pigs Fly, Inc. (York, ME).

cows' milk was significantly higher in the winter (116 \pm 23.1 $\mu g/250 \text{ ml}$) than in the summer (91.3 ± 16.6 $\mu g/250 \text{ ml}$, P =0.0004). Values for infant formulae ranged from 16.2–56.8 μ g iodine per 5 oz serving (mean, $23.5 \pm 13.78 \,\mu\text{g}/5 \,\text{oz}$), which was often far higher than labeled (Table 3).

Discussion

Recent reductions in U.S. dietary iodine intake have been ascribed to a possible reduction in iodine content in dairy products, to the removal of iodate conditioners in storebought breads, to new recommendations for reduced salt intake for blood pressure control, and to the increasing use of noniodized salt in manufactured or pre-made convenience foods (3, 5).

Inadequate dietary iodine is a significant worldwide health problem. Disorders caused by inadequate dietary iodine include goiter, subclinical or overt hypothyroidism, mental retardation, decreased fertility, increased neonatal mortality, and cretinism (6-8). Adequate iodine is particularly important for neural development in utero and in early life (9), and iodine deficiency remains the leading preventable cause of mental retardation worldwide (6). Furthermore, iodine deficiency will result in increased uptake of radioactive iodine and could, at least partially, explain the high prevalence of childhood thyroid cancer after the Chernobyl accident (10). Although chronic ingestion of excess iodine generally does not represent a significant public health problem, it may result in hypo- or hyperthyroidism in susceptible individuals, especially those with Hashimoto's thyroiditis and nodular nontoxic goiter, respectively. Excess iodine ingestion will also decrease the thyroid radioactive iodine uptake and reduce the effectiveness of radioactive iodine treatment for thyrotoxicosis and differentiated thyroid cancer.

Assessment of U.S. dietary iodine sources has not been performed for several years. Kidd et al. (11) obtained dietary frequency surveys and urinary iodine measurements from a sample of 754 schoolchildren between 1971 and 1972. Bread made with iodate conditioners, milk, and iodized salt were the primary sources of dietary iodine in their sample. National market basket samples performed for the Food and Drug Administration's Total Diet Study between 1982 and 1991 estimated that the average diet at the time contained iodine levels in excess of the current recommended daily allowance (3, 12).

Conditioners are added to store-bought bread to maintain

TABLE 3. Iodine content of infant formulae

Labeled iodine content (µg/5 oz)	Actual iodine content (µg/5 oz)
10	17.9
15	22.2
15	17.9
15	17.3
8	56.8
8	16.2
6	24.2
15	15.9
	10 15 15 15 8 8 6

- ^a Products of Mead Johnson Nutritionals (Evansville, IN).
- ^b Products of Nestle USA, Inc. (Glendale, CA).
- ^c Products of Ross Products (Columbus, OH).

freshness and prolong shelf life. In the 1960s, iodate bread conditioners were widely used. London et al. (13), in 1965, reported that bread was a source of large quantities of dietary iodine, with iodine content as high as 150 μg per slice. This was considered to be a contributing cause to the decreasing radioactive iodine uptake in the U.S. during the 1960s (14, 15). Because of the concerns about high bread iodine content, commercial bakeries now less commonly use iodate bread conditioners. The decreasing use of iodate bread conditioners is thought to have contributed to the reduction in dietary iodine levels between the 1970s and the early 1990s.

Three brands of bread in our study had inordinately high iodine content, up to 587 μ g iodine per slice, which was likely due to the presence of iodate bread conditioners. Thus, if an individual eats two slices of these breads over a 24-h period, up to 1,174 μg iodine will be ingested from bread alone, resulting in excess iodine exposure assuming an otherwise normal diet. The tolerable upper limit for daily iodine intake in adults established by the U.S. Institute of Medicine is 1,100 μ g (1). The ingredient list for only one of these breads indicated the use of an iodate conditioner. By contrast, the iodine content of several bread brands was low despite ingredient labeling that indicated use of iodate. It is clear that current labeling of bread does not accurately predict the content of iodine. Bread manufacturers should be encouraged to avoid the addition of additives containing excess iodine and to accurately list all iodine-containing substances and the iodine content per bread slice on an easily readable label.

Cows' milk continues to be a primary source of U.S. dietary iodine. Milk iodine content increased by 300-500% over the period from 1965–1980, largely due to changes in cattle feeds (16). The limitation of the allowable amount of organic iodine ethylenediamine dihydroiodine in cattle feed to 10 mg per cow daily in 1986 has resulted in decreases in the iodine content of U.S. cows' milk. Iodine supplementation of cattle feed may contribute to the iodine content of cows' milk in New England, but unlike seasonal variations in milk iodine content in the United Kingdom (17), the seasonal variation in iodine content of cows' milk noted in this study cannot be ascribed to alterations in the diet of dairy cows (Hines, J. G., Director, Commonwealth of Massachusetts Department of Food and Agriculture Division of Diary Services, personal communication, January 2003). Iodine is also introduced into cows' milk by the use of iodophor disinfectant in pre- and postmilking teat dips and udder washes. Such teat dips contain up to 1% available iodine and have been shown to significantly increase milk iodine residues by absorption through the skin and subsequent incorporation into milk (18, 19). Povidone iodine disinfecting solutions are also used to clean tanker trucks, vats, and milking equipment; however, by Food and Drug Administration regulations, such solutions may contain only 12.5–25 parts iodine per million (20), and residues from these solutions are unlikely to add a substantial amount of iodine to the milk supply (Hines, J. G., personal communication). The average iodine content of milk in our sample was 110 μ g per cup. The recommended intake for adequate calcium nutrition of 4 cups daily would provide 440 μ g iodine daily. This intake is 4.8 times the recommended intake for children and 2.9 times the recommended intake for adults. A combination of 4 cups of milk and two slices of bread could result in a daily ingestion of greater than 1,700 μg iodine, or 11.3 times the recommendations and 600 μg above adult tolerable limits. Generally, excess iodine intake is not a risk for thyroid dysfunction in the normal thyroids of children, but hypothyroidism or hyperthyroidism may occur in adults with preexisting Hashimoto's thyroiditis or nodular goiter, respectively. In view of the high iodine content in milk, it seems advisable that the iodine content should be incorporated into package labeling. Finally, infant formulae contain sufficient iodine, often more than labeled. They should provide infants with adequate iodine intake during this critical period of brain development.

Dietary iodine deficiency is an important public health problem worldwide. Overall, our findings, although based on small and nonrandom samples, suggest that U.S. iodine nutrition may remain an area worthy of public health concern. It is clear that there is a wide amount of variation in the iodine content of some common foods. Iodine content of foods is not well reflected by package labeling. There needs to be increased awareness of the importance of adequate iodine nutrition, particularly during pregnancy and lactation, among the U.S. public. Accurately identifying iodine content on food package labels would facilitate this process. Women of childbearing age should be encouraged to use iodine-containing multivitamins. Currently, many prenatal vitamins and other multivitamin preparations do not contain iodine (21). Salt iodization has been the mainstay of iodine deficiency prevention programs in countries around the world, and use of iodized salt should continue to be promoted in the United States. Finally, there is a need for larger and more systematic studies of iodine nutrition in different U.S. populations and for routine monitoring of food iodine content.

Acknowledgments

Received November 19, 2003. Accepted March 29, 2004.

Address all correspondence and requests for reprints to: Lewis E. Braverman, M.D., 88 East Newton Street, Evans 201, Boston, Massachusetts 02118. E-mail: lewis.braverman@bmc.org.

References

- Food and Nutrition Board Institute of Medicine 2001 Dietary reference intakes. Washington, DC: National Academy Press
- Hollowell JG, Staehling NW, Hannon WH, Flanders DW, Gunter EW, Maberly GF, Braverman LE, Pino S, Miller DT, Garbe PL, DeLozier DM, Jackson RJ 1998 Iodine nutrition in the United States. Trends and public health implications: iodine excretion data from National Health and Nutrition Examination Surveys I and III (1971–1974 and 1988–1994). J Clin Endocrinol Metab 83:3401–3408
- 3. **Pennington JA, Schoen SA** 1996 Total diet study: estimated dietary intakes of nutritional elements, 1982–1991. Int J Vitam Nutr Res 66:350–362
- Benotti J, Benotti N, Pino S, Gardyna H 1965 Determination of total iodine in urine, stool, diets and tissue. Clin Chem 11:932–936
- Lee K, Bradley R, Dwyer J, Lee SL 1999 Too much versus too little: the implications of current iodine intake in the United States. Nutr Rev 57:177–181
- Delange F 2000 Iodine deficiency. In: Braverman LE, Utiger RD, eds. Werner and Ingbar's the thyroid. 8th ed. Philadelphia: Lippincott Williams & Wilkins; 295–316
- Boyages SC 1993 Clinical review 49: iodine deficiency disorders. J Clin Endocrinol Metab 77:587–591
- Dunn JT 2003 Iodine deficiency and its elimination by iodine supplementation. In: Braverman LE, ed. Diseases of the thyroid. 2nd ed. Totowa, NJ: Humana Press; 329–345
- Glinoer D 2003 Feto-maternal repercussions of iodine deficiency during pregnancy. An update. Ann Endocrinol 64:37–44
- Jackson RJ, DeLozier DM, Gerasimov G, Borisova O, Garbe PL, Goultchenko L, Shakarishvili G, Hollowell JG, Miller DT 2002 Chernobyl and iodine deficiency in the Russian Federation: an environmental disaster leading to a public health opportunity. J Public Health Policy 23:453–470
- Kidd PS, Trowbridge FL, Goldsby JB, Nichaman MZ 1974 Sources of dietary iodine. J Am Diet Assoc 65:420–422
- Park YK, Harland BF, Vanderveen JE, Shank FR, Prosky L 1981 Estimation of dietary iodine intake of Americans in recent years. J Am Diet Assoc 79:17–24
- London WT, Vought VL, Brown F 1966 Bread: a dietary source of large quantities of iodine. N Engl J Med 223:338
- Sachs BA, Siegel E, Horwitt BN 1972 Bread iodine content and thyroid radioiodine uptake: a tale of two cities. Br Med J 1:79–81
- Pittman JA, Dailey GE, Beschi RJ 1969 Changing normal values for thyroidal radioiodine uptake. N Engl J Med 280:1431–1434
- Hemken RW 1980 Milk and meat iodine content: relation to human health. I Am Vet Med Assoc 176:1119–1121
- United Kingdom Ministry of Agriculture, Fisheries and Food 2000 Food surveillance sheet 198: iodine in milk. London: United Kingdom Ministry of Agriculture, Fisheries and Food
- Galton DM, Petersson LG, Erb HN 1986 Milk iodine residues in herds practicing iodophor premilking teat disinfection. J Dairy Sci 69:267–271
- Conrad LM, Hemken RW 1978 Milk iodine as influenced by an iodophor teat dip. J Dairy Sci 61:776–780
- 20. U.S. Code of Federal Regulations 21CFR178.1010
- 21. Lee SL, Roper J 2004 Inadequate iodine supplementation in American multivitamins. Endocr Pract 10(Suppl 1):46 (Abstract)

JCEM is published monthly by The Endocrine Society (http://www.endo-society.org), the foremost professional society serving the endocrine community.